MICROSATELLITE LETTERS

Development and characterization of microsatellite loci for two species of Beringian birds, rock sandpiper (*Calidris ptilocnemis*) and Pacific wren (*Troglodytes pacificus*)

Christin L. Pruett · Cesili Whelan · Angela Ricono · Stacey L. Lance · Travis Glenn · Brant Faircloth · Kevin Winker

Received: 7 August 2013 / Accepted: 17 September 2013 © Springer Science+Business Media Dordrecht 2013

Abstract Identification and assessment of small, endemic populations are priorities for conservation. We isolated and characterized 8 microsatellite loci from rock sandpiper (*Calidris ptilocnemis*) and 5 microsatellite loci from Pacific wren (*Troglodytes pacificus*), species with endemic populations of named subspecies that are of conservation concern. Eighteen to 20 individuals of each species from several locations in Alaska were screened for polymorphism. Loci for each species showed high polymorphism, with rock sandpiper ranging from 5 to 14 alleles per locus and 0.73–0.88 expected heterozygosity and Pacific wren ranging from 5 to 14 alleles per locus and 0.55–0.91 expected heterozygosity. Loci developed for rock sandpipers were also polymorphic in closely related taxa. These loci are the first developed for either species and will be

Electronic supplementary material The online version of this article (doi:10.1007/s12686-013-0040-4) contains supplementary material, which is available to authorized users.

C. L. Pruett (⊠) · C. Whelan · A. Ricono Department of Biological Sciences, Florida Institute of Technology, Melbourne, FL 32901, USA e-mail: cpruett@fit.edu

S. L. Lance Savannah River Ecology Laboratory, University of Georgia, Aiken, SC 29802, USA

T. Glenn Department of Environmental Health Science, University of Georgia, Athens, GA 30602, USA

B. Faircloth Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095, USA

K. Winker University of Alaska Museum, Fairbanks, AK 99775, USA used to identify and conserve endemic populations in the Bering Sea region.

Keywords *Troglodytes* · *Calidris* · PCR primers · Microsatellite · Aleutian Islands · Pribilof Islands

Rock sandpiper (*Calidris ptilocnemis*) and Pacific wren (*Troglodytes pacificus*) exhibit high levels of intraspecific diversity within Beringia, with endemic populations (named subspecies) primarily found on remote oceanic islands including Pribilof and Aleutian island populations that are of conservation concern (Gotthardt et al. 2012). However, little is known about the genetic diversity or connectivity among populations. Highly polymorphic genetic markers are needed to assess population structure and gene flow and to identify isolated endemic populations that should be treated as independent management units.

Total genomic DNA was extracted from tissue samples of two individuals for each species from Alaska populations. Microsatellite loci were isolated following Glenn and Schable (2005). Briefly, DNA was digested with restriction enzymes, adapters were ligated onto ends of fragments, biotinylated probes and streptavidin beads were used to physically separate microsatellite-containing DNA fragments, and PCR was done to increase the yield of fragments. DNA fragments were then run on a Roche 454 FLX with Titanium chemistry along with similarly treated libraries, each with unique adapter and Roche MID-tag combinations. We de-multiplexed with demuxipy (https:// github.com/faircloth-lab/demuxipy/) and used a customized version (https://github.com/faircloth-lab/msatcommandergs) of MSATCOMMANDER (Faircloth 2008) to find 10,247 and 15,954 sequences containing microsatellites from 15,701 to 26,433 sequence reads of sandpiper and

Table 1 Ir	formation about 8 microsatellite loci developed for rock sandpiper (Calidri	ptilocnemis; Capt) and 5 mic	rosatellite loci fo	r Pacific v	vren (Trog	glodytes pac	ificus; Trpa)	
Locus	Primer sequence 5'-3'	Repeat motif	Size (bp)	Ν	А	H_o	H_e	TD
Capt4	F: CAGTCGGGCGTCATCATTTGTCCACCAGGACTTAGAAA R: CTCCCAAGGCAGGCTGAAA	(AGAT) ¹²	206–226	16	9	0.81	0.78	55/45
Capt6	F: CAGTCGGGCGTCATCACATACTGGAAGGCCGCTA R: CCTCGCTCCAACATCATGC	(ATCT) ¹⁰	158–194	18	6	0.94	0.84	60/50
Capt14	F: CAGTCGGGCGTCATCACTGTCTGAAACATATTGGCAAGC R: TGAAGAGATCCAGAAGATCCGT	(ATCT) ¹²	360–396	17	11	0.65	0.88*	55/45
Capt16	F: GGTGTGACAACATCATGGCAG R: CAGTCGGGGGGTCATCAAACTGGACAGGCATCCAAA	(ATCT) ¹²	203–227	17	S	0.47	0.73	65/55
Capt18	F: CAGTCGGGCGTCATCAAACCTGCCACCTGATGGAT R: TCTTACCTATGTGTGGTCAAGC	(AGAT) ⁹	288–312	16	Γ	0.75	0.81	55/45
Capt20	F: CAGTCGGGCGTCATCATACTGGACCACTTGTGGCAT R: TCTACTGCATCCTGTTTCCCA	(AGAT) ⁸	328–360	17	14	0.47	0.88*	55/45
Capt23	F: CAGTCGGGCGTCATCACTTCACATACTGGAAGGCCG R: CCTCGCTCCAACATCATGC	(ATCT) ¹⁰	163–199	16	6	1.00	0.83	60/50
Capt43	F: TGGTGTGGTAGCAATCTCCT R: CAGTCGGGCGTCATCACTGCCATCCAACAAGATGCC	(GGAT) ¹⁴	269–305	18	6	0.56	0.81*	55/45
Trpa4	F: CAGTCGGGCGTCATCAAGGTCGTTCAGGCTTGAGT R: CTCTACACTCTACAACCGTTCC	(AC) ¹³	314–326	20	Γ	0.50	0.69	60/50
Trpa12	F: CAGTCGGGCGTCATCATATCAGCACTGTTCCAGCCA R: TGAAATACAGTAGGGAACCAAGTC	(ACT) ¹⁶	220–253	20	17	09.0	0.79*	55/45
Trpa21	F: CTCCCAGGCATCCCTCATTC R: CAGTCGGGCGTCATCACCAAGAGTCAGCTTCCACA	(AGAT) ¹⁴ (AGAT) ⁹	328–396	20	14	0.65	0.91	60/50
Trpa22	F: CAGTCGGGCGTCATCAGCCAACAGTAGGACATGGCT R: CAGTGTTAAGGTTGCTACACCA	(GT) ¹⁰	210–218	19	S	0.16	0.55*	60/50
Trpa27	F: CAGTCGGGCGTCATCACTCACTCTGGCCCAATTCCT R: GAGAACTTGGCTCTTCATCA	(AC) ¹⁰	227–241	20	Ś	0.50	0.60	65/55

Size indicates range of alleles in base-pairs (bp) including CAG tag, number of individuals genotyped (N), number of alleles (A), observed heterozygosity (H_o), expected heterozygosity (H_e), and TD refers to the annealing temperatures for the touchdown protocol for each locus. Loci out of Hardy–Weinberg equilibrium (P < 0.01) are indicated with an asterisk next to expected heterozygosity values

Author's personal copy

wren, respectively. MSATCOMMANDER was used to design primers with the CAG primer sequence (5'-CAG-TCGGGCGTCATCA-3') prepended to the 5' end of one primer to enable use of a third fluorescently labeled primer identical to the CAG sequence during PCR (Hauswaldt and Glenn 2003).

Forty-five primer pairs for rock sandpiper and 28 primer pairs for Pacific wren were tested for amplification and polymorphism using genomic DNA from 5 individuals for each species. Amplifications were in 20 ul volumes (250ug/ mL BSA, 10×Buffer B (Fisher Scientific), 25 mM MgCl₂, 5 uM unlabeled primer, 0.5 uM tag labeled primer, 5 uM universal dye-labeled primer, 2.5 mM dNTPs, 0.5 units Taq DNA polymerase (Fisher Scientific), and 20 ng DNA template) using a BioRad MyCycler or Multigene Labnet International thermal cycler. We used touchdown cycling conditions to amplify DNA and to attach the universal dyelabeled primer. Parameters consisted of an initial denaturation step of 2 min 30 s at 95° C then 20 cycles of 95° C for 20 s, 65°-50° C annealing temperature for 20 s (decreasing 0.5° C per cycle), and extension step of 72° C for 30 s followed by 15 cycles of 95° C for 20 s, 55°–45° C for 20 s, and 72° C for 30 s. Cycles were followed with a final extension step of 72° C for 10 min. Amplifications were run on an ABI3730XL sequencer (Applied Biosystems).

We examined polymorphic loci using 18 individuals for rock sandpiper from several Alaska locations, including Cold Bay, Nome, Adak Island, Shemya Island, the Pribilof Islands, and Kodiak Island, and two individuals each for three closely related species, purple sandpiper (Calidris maritima), dunlin (Calidris alpina), and sanderling (Calidris alba). Twenty Pacific wrens were genotyped from Juneau, Near Islands, Rat Islands, and Middleton Island in Alaska. Eight of the rock sandpiper and 5 of the Pacific wren primers amplified DNA and exhibited polymorphism (Table 1, Online Resource 1). We scored alleles using GeneMapper software (Applied Biosystems) and evaluated the number of alleles per locus (A), observed and expected heterozygosity (H_0 and H_e), Hardy–Weinberg equilibrium, and linkage equilibrium using Arlequin ver 3.5 (Excoffier et al. 2005). We found that the number of alleles per locus

ranged from 5 to 14 for both species. Observed heterozygosity for rock sandpiper ranged from 0.47 to 1.00 and for Pacific wren from 0.16 to 0.65, and expected heterozygosity ranged from 0.73 to 0.88 for rock sandpiper and from 0.55 to 0.91 for Pacific wren. All loci were in linkage equilibrium but several loci were out of Hardy–Weinberg equilibrium after correction for multiple tests (Table 1). Deviations from Hardy–Weinberg are expected given that samples were taken from several populations that are likely to have dissimilar histories.

All 8 loci that were polymorphic in rock sandpipers were successfully amplified in purple sandpiper, dunlin, and sanderling. Most of these were polymorphic for each species, except for Capt16 for purple sandpiper and dunlin, Capt 4 and Capt 18 for sanderling, and Capt20 for dunlin. However, these loci might be polymorphic if a larger sample size was assessed. We suggest that these loci could be successfully used for population-level assessments in these species and possibly in other members of the genus *Calidris*.

Acknowledgments This work was supported by the Florida Institute of Technology, University of Alaska Museum, and by the U. S. Department of Energy under Award Number DE–FC09– 07SR22506 to the University of Georgia Research Foundation. We thank M. Zimmerman, S. Garcia, M. Smith, and R. Selvam for help in the laboratory.

References

- Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. Evol Bioinforma Online 1:47–50
- Faircloth BC (2008) MSATCOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. Mol Ecol Resour 8:92–94
- Glenn TC, Schable NA (2005) Isolating microsatellite DNA loci. Methods Enzymol 395:202–222
- Gotthardt TA, Walton KM, Fields TL (2012) Setting priorities for Alaska's Wildlife Action Plan. University of Alaska Anchorage, Alaska Natural Heritage Program
- Hauswaldt JS, Glenn TC (2003) Microsatellite DNA loci for the diamondback terrapin (*Malaclemys terrapin*). Mol Ecol Resour 3:174–176